

The 24 μ m Luminosity Function of spectroscopic SWIRE sources from the Lockman Validation Field.

N.Onyett¹, S.Oliver¹, G.Morrison⁴, F.Owen², F.Pozzi³, D.Carson¹,
 SWIRE team⁴

¹*Astronomy Centre, University of Sussex, Brighton, UK* ²*NRAO, Socorro, USA* ³*Dipartimento Astronomia Bologna, Italy* ⁴*IPAC, CalTech, Pasadena, CA, USA*

Abstract. A spectroscopic follow-up of SWIRE sources from the Lockman Validation Field has allowed the determination of the SWIRE/WIYN 24 μ m Luminosity Function (LF). The spectroscopic sample was chosen above a 24 μ m flux limit at 260 μ Jy and an r-band optical limit of $r < 21$. A spectroscopic completeness of 82.5% was achieved. We found the median redshift for the sample to be $z_{\text{med}} = 0.29$. Markov-Chain Monte-Carlo (MCMC) and Maximum Likelihood Estimator (MLE) techniques were employed to fit a parametric LF. Our result of the local LF (LLF) is consistent with the local 25 μ m determination from Shupe et al. (1998). We split the sample at a redshift of $z_{\text{split}} = 0.36$ and find strong evidence for galaxy evolution.

1. Introduction

Since one can not study directly how a galaxy forms and evolves (how the mass and luminosity of a galaxy change with time), it becomes necessary to analyse the statistics of a large number of galaxies at different epochs. Using the SWIRE (Lonsdale et al. 2004) 24 μ m MIPS-band, one is able observe the FIR emission from dust, tracing star-formation (SF) regions and SF-galaxies. By studying these galaxies at different epochs one can gain an estimate of the star formation history of the Universe. As a follow up to SWIRE, a spectroscopic redshift survey covering a 0.55 deg² region, centred in the Lockman Validation Field was undertaken. The sample was observed using the HYDRA spectrometer on the 4-m WIYN telescope at KPNO and included 24 μ m sources down to a Vega r-band magnitude of $r < 21$. A total of 286 24 μ m sources were observed and redshifts for the resultant spectra were determined. Redshifts for 170 sources with 24 μ m fluxes down to a flux limit of 260 μ Jy were determined, giving a spectroscopic completeness for the sample of 82.5%.

2. Constructing SWIRE/WIYN LF₂₄

In order to determine each object's 24 μ m luminosity (L_{24}) an appropriate SED and K-correction for that galaxy must be chosen. Here 6 SEDs were chosen, 2 AGN, 2 spiral and 2 starburst (Xu et al., 2001). The fitting was performed using a minimum χ^2 technique that incorporated an error weighting (including a dispersion factor, previously unseen in the literature). This dispersion factor




Figure 1. High [pale-line] & low [dark-line] z-sample LF_{24} , split at $z=0.36$. Note that incompleteness corrections have not been made in faintest bins.




Figure 2. Left: MCMC & MLE best fits Right: comparison to Shupe's LF_{25}

accounts for the scatter within a particular population of galaxies in colour-colour space. Each SED was then convolved through the $24\mu\text{m}$ filter. The LF_{24} was estimated by implementing the $1/V_{\text{max}}$ estimator technique (Schmidt 1968) and took into account both the r-band and $24\mu\text{m}$ flux limits. A redshift split at $z_{\text{split}} = 0.36$ was imposed, allowing evolution in the sample to be investigated, see Fig.1. The high-z sample (pale line) is composed of 38% of the total z-sample with $V/V_{\text{max}} = 0.72 \pm 0.04$, showing strong evidence for luminosity evolution. The low-z sample (dark line) is composed of 62% of the total z-sample with $V/V_{\text{max}} = 0.46 \pm 0.03$. Note: incompleteness corrections need to be applied to the faintest luminosity bins of the high-z sample. Parameterization of the SWIRE LF was performed and a comparison with low-z from IRAS was made. Using MCMC (Kashyap 1998) & MLE (Sandage, et al. 1979) techniques to fit Saunders et al. 1990 parametric form of the LF, the best fitting MCMC and MLE model fits to the SWIRE LF_{24} are shown in Fig.2. A direct comparison to Shupe et al. 1998 LF_{25} is also shown in Fig.2. Since this is an apriori model and not a fit it provides a fairly good description of our data.

References

- Kashyap et al. 1998, ASPC, 154, 844 Schmidt.M 1968, ApJ, 151, 393
 Lonsdale et al. 2004, ApJ, 1, 54 Shupe et al. 1998, ApJ, 501, 597
 Sandage, et al. 1979, ApJ, 232, 352 Xu et al. 2001, ApJ, 562, 179
 Saunders et al. 1990, MNRAS, 242, 318